


BioMag[®] SelectaPure[™] Anti-Human CD2 Antibody Particles

Description

The CD2 antigen, also known as leukocyte function-associated antigen 2 (LFA-2), is a monomeric 50 kDa glycoprotein that is a receptor for sheep erythrocytes and the CD58/LFA-3 ligand. CD2 plays a role in T-cell signaling and lymphocyte adhesion. It is expressed on the vast majority of thymocytes and all peripheral T lymphocytes and is also expressed on malignant cells of T-cell origin including T-ALL and a subset of NK cells.

BioMag Anti-Human CD2 Antibody Particles are designed for positive selection of CD2 plus human T-cells.



General Recommendation*:

Conc. #	4×10^8 particles/ml
Vol. Used	50 μ l
# Particles	2×10^7 per test
# Target Cells	Approximately 7.32×10^5
Particles:Cell Ratio	27:1
% Depletion	94%

*These values should be used as a starting point in optimizing experimental protocols. Due to differences in the distribution of cell types in samples and other variables, the researcher is strongly encouraged to determine the optimal particle to cell ratios for their experiments.

Cell sorting results using BioMag Selectapure anti-human CD2 leukocyte particles for positive selection. Typically whole blood or purified leukocytes and particles are incubated for 30 minutes at room temperature and then magnetically separated. The supernatant is collected, incubated with the appropriate two-color antibody cocktail, and then analyzed by flow cytometry. Figure A depicts the cell population prior to positive selection. Figure B depicts the cell population after positive selection. The particle to cell ratios reported above are based on experiments where cells were exposed to the particles once.

Particle Concentration

The concentration of BioMag Anti-Human CD2 is approximately 4.0 mg/ml. There are approximately 1×10^8 BioMag particles per mg.

Cell Separation Recommendations

Depending upon antigen availability and the size of the target cell population, cell sorting applications may require up to 50-60 magnetic particles per cell based on the target cell population. Magnetic particles and cells should be incubated at room temperature for 30 minutes to one hour in media containing 5-10% protein (to reduce non-specific binding) for successful separation. (Note: Increasing the incubation time beyond one hour may be necessary to achieve the desired depletion.) Each researcher must optimize particle to cell ratio and incubation time for their application.

Some applications require the detachment of BioMag antibody particles from cells after separation. One approach would involve culturing cells after positive selection. Cultures can be maintained for about 48 hours during which magnetic particles fall away from cells due to cell surface changeover. The magnetic particles are then easily removed via a magnetic separation. Another approach is the use of a protease such as chymopapain to break the antigen-antibody bond and remove the particles magnetically. Depending upon the application, it may not be necessary to remove the cells from the BioMag particles. BioMag particles are only 1 μ m in size and have been successfully used in FACS equipment. They will not jam the machine and are distinguishable from cells. Alternatively, negative selection approaches can be very effective in producing specific cell populations.

Storage and Stability

The suspension is supplied in PBS/EDTA/1.0% BSA/0.1% sodium azide buffer at pH 7.5. Washing BioMag Anti-Human CD2 particles in sterile media to remove preservative prior to use is recommended. Using a magnetic separation unit for washing instead of centrifugation is strongly recommended. Do not freeze, dry or centrifuge BioMag particles. Freezing, drying and centrifuging BioMag particles can result in aggregation and loss of binding activity. BioMag Anti-Human CD2 particles are stable when stored at 4°C.

Safety

BioMag Anti-Human CD2 particle suspension contains sodium azide. Sodium azide may react with lead and copper plumbing to form explosive metal azides. Upon disposal of material, flush with a large volume of water to prevent azide accumulation. Please consult the Material Safety Data Sheet for more information.

Ordering Information:

Cat. #	Description	Size
85002	BioMag Anti-Human CD2	1 ml
	Antibody Particles	5 ml

To Order:

In The U.S. Call: 1-800-523-2575 • 215-343-6484

In The U.S. FAX: 1-800-343-3291 • 215-343-0214

In Germany Call: (49) 6221-765767

In Germany FAX: (49) 6221-764620

Order online anytime at www.polysciences.com

This product is for research use only and is not intended for use in humans or for in vitro diagnostic use.

Should any of our materials fail to perform to our specifications, we will be pleased to provide replacements or return the purchase price. We solicit your inquiries concerning all needs for life sciences work. The information given in this bulletin is to the best of our knowledge accurate, but no warranty is expressed or implied. It is the user's responsibility to determine the suitability for his own use of the products described herein, and since conditions of use are beyond our control, we disclaim all liability with respect to the use of any material supplied by us. Nothing contained herein shall be construed as a recommendation to use any product or to practice any process in violation of any law or any government regulation.