

U.S. Corporate Headquarters 400 Valley Rd. Warrington, PA 18976 1(800) 523-2575 / (215) 343-6484 1(800)343-3291 fax info@oolysciences.com Polysciences Europe GmbH Badener Str. 13 69493 Hirschberg an der Bergstrasse, Germany +(49) 06201-845200 +(49) 06201-8452020 fax

info@polysciences.de

2F-1, 207 DunHua N. Rd. Taipei, Taiwan 10595 (886) 2 8712 0600 (886) 2 8712 2677 fax info@polysciences.tw

Polysciences Asia-Pacific, Inc.

TECHNICAL DATA SHEET 594

Page 1 of 2

BioMag® SelectaPure anti-Mouse CD45R

Catalog Number: 85145

DESCRIPTION

The B220 isoform of the CD45 antigen is a subset of the CD45 glycoprotein family that is expressed on all hematopoietic cells with the exception of mature erythrocytes.

BioMag® SelectaPure anti-Mouse CD45R particles are designed for positive selection of hematopoietic cells with the exception of the mature erythrocytes. The antibody used reacts with cells expressing a CD45R (B220) phenotype, primary cells of B lineage, including pro B cells through mature B cells. The antibody can also react with a subset of activated T cells and with non-MHC-restricted active lymphokine-activated killer cells.

BioMag® SelectaPure anti-Mouse CD45R is a suspension of magnetic particles approximately 1.5µm in size. The suspension is supplied in a phosphate buffered saline (pH 7.5) containing EDTA, 1.0% BSA, and 0.1% sodium azide.

CHARACTERISTICS

Mean Diameter: ~1.5μm Particle Concentration: 1 mg/ml

Particle Count: 1 x 10⁸ BioMag[®] particles per mg

PROCEDURE


Researchers are advised to optimize the use of particles in any application.


Depending on antigen availability and the size of the target cell population, cell sorting applications may require up to 50-60 magnetic particles per cell. Magnetic particles and cells should be incubated at room temperature for 30-60 minutes in media containing 5-10% protein (to reduce nonspecific binding) for successful separation. Gentle end-over-end mixing or rocking during incubation is required for optimal results. (*Note*: Increasing the incubation time beyond one hour may be necessary to achieve the desired depletion.)

Some applications require the detachment of BioMag® antibody particles from cells after separation. One approach would involve culturing cells after positive selection. Cultures can be maintained for about 48 hours during which magnetic particles fall away from cells due to cell surface changeover. The magnetic particles are then easily removed via a magnetic separation. Another approach is the use of a protease, such as chymopapain, to break the antigen-antibody bond and remove the particles magnetically. Depending upon the application, it may not be necessary to remove the cells from the BioMag® particles. BioMag® particles are only 1-2µm in size and have been successfully used in FACS equipment. They

will not jam the machine and are distinguishable from cells. Alternatively, negative selection approaches can be very effective in producing specific cell populations.

Cell sorting results using BioMag® SelectaPure anti-Mouse CD45R leukocyte particles for positive selection. Typically, whole blood or purified leukocytes and particles are incubated for 30 minutes at room temperature and then magnetically separated. The supernatant is collected, incubated with the appropriate two-color antibody cocktail, and then analyzed by flow cytometry. Figure A depicts the cell population prior to positive selection. Figure B depicts the cell population after positive selection. The particle to cell ratios reported above are based on experiments where cells were exposed to the particles once.





General Recommendation*:

Concentration# 1.00 x 10⁸ particles/ml

Volume Used 0.025ml

Particles 2.5 x 10 6 per mL of blood

Target Cells 2.5 x 10^5 per ml Particle:Target Cell Ratio 11:1

% Depletion 97.0%

STORAGE AND SAFETY

Storage Store at 4°C. Freezing, drying, or centrifuging BioMag® may result in irreversible aggregation and loss of binding activity. Washing BioMag® SelectaPure anti-Mouse CD45R particles in sterile media to remove preservative prior to use is recommended. Using a magnetic separation unit for washing instead of centrifugation is also strongly recommended.

^{*} These values should be used as a starting point in optimizing experimental protocols. Due to differences in the distribution of cell types in samples and other variables, the researcher is strongly encouraged to determine the optimal particle to cell ratios for their experiments.

TECHNICAL DATA SHEET 594

Page 2 of 2

Safety This particle suspension contains sodium azide. Sodium azide may react with lead and copper plumbing to form explosive metal azides. Upon disposal of material, flush with a large volume of water to prevent azide accumulation. Please consult the Safety Data Sheet for more information.

These products are for research use only and are not intended for use in humans or for *in vitro* diagnostic use.

ORDERING INFORMATION

Cat. #	Description	Size
85145-5	BioMag® SelectaPure anti-Mouse CD45R	5ml

TO ORDER

In The U.S. Call: 1(800) 523-2575 • (215) 343-6484 In The U.S. Fax: 1(800) 343-3291 • (215) 343-0214

In Germany Call: +(49) 06201-845200 In Germany Fax: +(49) 06201-8452020 In Asia Call: (886) 2 8712 0600

In Asia Fax: (886) 2 8712 2677

Order online anytime at www.polysciences.com