



U.S. Corporate Headquarters  
400 Valley Rd.  
Warrington, PA 18976  
1(800) 523-2575 / (215) 343-6484  
1(800)343-3291 fax  
info@polysciences.com

Polysciences Europe GmbH  
Badener Str. 13  
69493 Hirschberg an der Bergstrasse,  
Germany  
+(49) 06201-845200  
+(49) 06201-8452020 fax  
info@polysciences.de

Polysciences Asia-Pacific, Inc.  
2F-1, 207 DunHua N. Rd.  
Taipei, Taiwan 10595  
(886) 2 8712 0600  
(886) 2 8712 2677 fax  
info@polysciences.tw

## TECHNICAL DATA SHEET 586

Page 1 of 2

# BioMag<sup>®</sup> SelectaPure anti-Human CD19

Catalog Number: 85019

### DESCRIPTION

The CD19 antigen is expressed on all normal B lymphocytes and pro-B lymphocytes. CD19 expression is lost as the B lymphocytes progress to mature plasma cells. It is also found on the surface of follicular dendritic cells, early cells of myelomonocytic lineage, and on stabilized B cell lines. It is not present on T lymphocytes, NK cells, monocytes, and granulocytes. The CD19 antigen is a 95 kDa type I integral membrane glycoprotein involved in the regulation of B lymphocyte development, activation, and differentiation.

BioMag<sup>®</sup> SelectaPure anti-Human CD19 particles are used for positive selection of B lymphocytes. BioMag<sup>®</sup> SelectaPure anti-Human CD19 particles recognize most B cell lines and react with the great majority of non-T acute lymphocytic leukemias. Research applications for this product include leukemia typing, identification of B cell lymphomas, and investigation of the ontogeny and differentiation of human B lymphocytes.

BioMag<sup>®</sup> SelectaPure anti-Human CD19 is a suspension of magnetic particles approximately 1.5 $\mu$ m in size. The suspension is supplied in a phosphate buffered saline (pH 7.5) containing EDTA, 1.0% BSA, and 0.1% sodium azide.

### CHARACTERISTICS

Mean Diameter: ~1.5 $\mu$ m  
Particle Concentration: 4mg/ml  
Particle Count: 1 x 10<sup>8</sup> BioMag<sup>®</sup> particles per mg

### PROCEDURE


Researchers are advised to optimize the use of particles in any application.

Depending on antigen availability and the size of the target cell population, cell sorting applications may require up to 50-60 magnetic particles per cell. Magnetic particles and cells should be incubated at room temperature for 30-60 minutes in media containing 5-10% protein (to reduce non-specific binding) for successful separation. Gentle end-over-end mixing or rocking during incubation is required for optimal results. (*Note:* Increasing the incubation time beyond one hour may be necessary to achieve the desired depletion.)

Some applications require the detachment of BioMag<sup>®</sup> antibody particles from cells after separation. One approach would involve culturing cells after positive selection. Cultures can be maintained for about 48 hours during which magnetic particles fall away from cells due to cell surface changeover. The magnetic particles are then easily removed via a magnetic separation. Another approach is the use of a protease, such as

chymopapain, to break the antigen-antibody bond and remove the particles magnetically. Depending upon the application, it may not be necessary to remove the cells from the BioMag<sup>®</sup> particles. BioMag<sup>®</sup> particles are only 1-2 $\mu$ m in size and have been successfully used in FACS equipment. They will not jam the machine and are distinguishable from cells. Alternatively, negative selection approaches can be very effective in producing specific cell populations.

**Cell sorting results using BioMag<sup>®</sup> SelectaPure anti-Human CD19 leukocyte particles for positive selection.** Typically, whole blood or purified leukocytes and particles are incubated for 30 minutes at room temperature and then magnetically separated. The supernatant is collected, incubated with the appropriate two-color antibody cocktail, and then analyzed by flow cytometry. Figure A depicts the cell population prior to positive selection. Figure B depicts the cell population after positive selection. The particle to cell ratios reported above are based on experiments where cells were exposed to the particles once.



#### General Recommendation\*:

Concentration#	4.0 x 10 <sup>8</sup> particles/ml
Volume Used	0.0125ml
# Particles	5.00 x 10 <sup>6</sup> per test
# Target Cells	1.35 x 10 <sup>5</sup> per test
Particle:Target Cell Ratio	37.0
% Depletion	99.22%

\* These values should be used as a starting point in optimizing experimental protocols. Due to differences in the distribution of cell types in samples and other variables, the researcher is strongly encouraged to determine the optimal particle to cell ratios for their experiments.

**STORAGE AND SAFETY**

**Storage** Store at 4°C. Freezing, drying, or centrifuging BioMag® may result in irreversible aggregation and loss of binding activity. Washing BioMag® SelectaPure anti-Human CD19 particles in sterile media to remove preservative prior to use is recommended. Using a magnetic separation unit for washing instead of centrifugation is also strongly recommended.

**Safety** This particle suspension contains sodium azide. Sodium azide may react with lead and copper plumbing to form explosive metal azides. Upon disposal of material, flush with a large volume of water to prevent azide accumulation. Please consult the Safety Data Sheet for more information.

**These products are for research use only and are not intended for use in humans or for *in vitro* diagnostic use.**

**ORDERING INFORMATION**

Cat. #	Description	Size
85019-5	BioMag® SelectaPure anti-Human CD19	5ml

**TO ORDER**

In The U.S. Call: 1(800) 523-2575 • (215) 343-6484  
 In The U.S. Fax: 1(800) 343-3291 • (215) 343-0214  
  
 In Germany Call: +(49) 06201-845200  
 In Germany Fax: +(49) 06201-8452020  
  
 In Asia Call: (886) 2 8712 0600  
 In Asia Fax: (886) 2 8712 2677

Order online anytime at [www.polysciences.com](http://www.polysciences.com)

Should any of our materials fail to perform to our specifications, we will be pleased to provide replacements or return the purchase price. We solicit your inquiries concerning all needs for life sciences work. The information given in this bulletin is to the best of our knowledge accurate, but no warranty is expressed or implied. It is the user's responsibility to determine the suitability for their own use of the products described herein, and since conditions of use are beyond our control, we disclaim all liability with respect to the use of any material supplied by us. Nothing contained herein shall be construed as a recommendation to use any product or to practice any process in violation of any law or any government regulation.